

## CDC's Advisory Committee on Immunization Practices Recommendations

The groups recommended to receive the 2009 H1N1 influenza vaccine include:

- **Pregnant women** because they are at higher risk of complications and can potentially provide protection to infants who cannot be vaccinated;
- **Household contacts and caregivers for children younger than 6 months of age** because younger infants are at higher risk of influenza-related complications and cannot be vaccinated. Vaccination of those in close contact with infants younger than 6 months old might help protect infants by “cocooning” them from the virus;
- **Healthcare and emergency medical services personnel** because infections among healthcare workers have been reported and this can be a potential source of infection for vulnerable patients. Also, increased absenteeism in this population could reduce healthcare system capacity;
- **All people from 6 months through 24 years of age**
  - **Children from 6 months through 18 years of age** because cases of 2009 H1N1 influenza have been seen in children who are in close contact with each other in school and day care settings, which increases the likelihood of disease spread, and
  - **Young adults 19 through 24 years of age** because many cases of 2009 H1N1 influenza have been seen in these healthy young adults and they often live, work, and study in close proximity, and they are a frequently mobile population; and,
- **Persons aged 25 through 64 years who have health conditions associated with higher risk of medical complications from influenza.**

No shortage of 2009 H1N1 vaccine is expected, but vaccine availability and demand can be unpredictable and there is some possibility that initially, the vaccine will be available in limited quantities. So, the ACIP also made recommendations regarding which people within the groups listed above should be prioritized if the vaccine is initially available in extremely limited quantities.

IF the H1N1 influenza vaccine is initially available in extremely limited quantities, the groups recommended to receive the 2009 H1N1 influenza vaccine include:

- Pregnant women,
- People who live with or care for children younger than 6 months of age,
- Health care and emergency medical services personnel with direct patient contact,
- Children 6 months through 4 years of age, and
- Children 5 through 18 years of age who have chronic medical conditions.

Once the demand for vaccine for the prioritized groups has been met at the local level, programs and providers should also begin vaccinating everyone from the ages of 25 through 64 years. Current studies indicate that the risk for infection among persons age 65 or older is less than the risk for younger age groups. However, once vaccine demand among younger age groups has been met, programs and providers should offer vaccination to people 65 or older.